



## Review

# Effective use of water (EUW) and not water-use efficiency (WUE) is the target of crop yield improvement under drought stress

A. Blum\*

Plantstress.com, P.O. Box 16246, Tel Aviv, Israel

## ARTICLE INFO

## Article history:

Received 12 December 2008

Received in revised form 19 March 2009

Accepted 19 March 2009

## Keywords:

Water-use efficiency

Drought resistance

Transpiration

Stomata

Grain yield

Harvest index

Plant breeding

Soil moisture

Roots

## ABSTRACT

Water-use efficiency (WUE) is often considered an important determinant of yield under stress and even as a component of crop drought resistance. It has been used to imply that rainfed plant production can be increased per unit water used, resulting in “more crop per drop”.

This opinionated review argues that selection for high WUE in breeding for water-limited conditions will most likely lead, under most conditions, to reduced yield and reduced drought resistance. As long as the biochemistry of photosynthesis cannot be improved genetically, greater genotypic transpiration efficiency (TE) and WUE are driven mainly by plant traits that reduce transpiration and crop water-use, processes which are crucially important for plant production. Since biomass production is tightly linked to transpiration, breeding for maximized soil moisture capture for transpiration is the most important target for yield improvement under drought stress. Effective use of water (EUW) implies maximal soil moisture capture for transpiration which also involves reduced non-stomatal transpiration and minimal water loss by soil evaporation. Even osmotic adjustment which is a major stress adaptive trait in crop plants is recognized as enhancing soil moisture capture and transpiration. High harvest index (HI) expresses successful plant reproduction and yield in terms of reproductive functions and assimilate partitioning towards reproduction. In most rainfed environments crop water deficit develops during the reproductive growth stage thus reducing HI. EUW by way of improving plant water status helps sustain assimilate partitions and reproductive success. It is concluded that EUW is a major target for yield improvement in water-limited environments. It is not a coincidence that EUW is an inverse acronym of WUE because very often high WUE is achieved at the expense of reduced EUW.

© 2009 Elsevier B.V. All rights reserved.

The coined slogan “more crop per drop” (Kijne et al., 2003) as a target for crop improvement in water-limited environments emerged in recent years in the press and among research administrators and sponsors. It is a very catchy slogan indeed, but also a misleading one. It does not serve well the cause of breeding for water-limited environments, especially rainfed conditions. It led breeders to believe that crop production under water-limited conditions can be genetically improved by increasing plant production per given amount of water used by the crop. A misconception also developed that improved water-use efficiency (WUE) is synonymous with drought resistance and high yield under drought stress. It is possible to achieve “more crop per drop” by certain crop and soil management practices (e.g. plant nutrition). However, this review concentrates on genetic improvement. It is not a comprehensive review of WUE.

This paper is therefore designed to clarify the meaning and consequences of WUE if used in breeding either in practice or in

concept. It raises the argument that the important determinant of plant production under most conditions of limited water supply is the effective use of water (EUW) and not high WUE.

Passioura (1996) proposed to view grain yield as a partial function of WUE:

$$Y = WU \times WUE \times HI \quad (1)$$

where  $Y$  is grain yield,  $WU$  is water-use, and  $HI$  is harvest index.

While Passioura shifted in opinion towards “water productivity” as a prime consideration in dryland crop production (Passioura, 2006), Eq. (1) remains quite popular among breeders (e.g. Reynolds and Tuberosa, 2008) and in training courses since it is simple and has some educational merit. Here it is used only as a gateway for explaining the main message of this review.

The equation implies that WUE is an independent variable in affecting grain yield. In this expression WUE equals  $B/WU$ , where  $B$  is biomass, therefore:

$$Y = WU \times \left( \frac{B}{WU} \right) \times HI \quad (2)$$

\* Tel.: +972 3 5270558; fax: +972 3 5246247.

E-mail address: [ablum@plantstress.com](mailto:ablum@plantstress.com).

In Eq. (2) WU can be cancelled out and then we return to the basics (Donald and Hamblin, 1976), namely:

$$Y = B \times HI \quad (3)$$

de Wit (1958) set the cornerstone for relating plant production to water-use:

$$B = \frac{mT}{E_0} \quad (4)$$

where  $B$  is crop biomass,  $m$  is a crop constant,  $T$  is crop transpiration and  $E_0$  is free water (potential) evaporation. This relationship is very solid and has stood the test of time. For genotypic comparisons  $E_0$  is common and can be removed from Eq. (4). Thus for genotypic comparisons, biomass of a genotype is a function of crop transpiration and a crop constant which is independent of  $T$ . A comparison of  $m$  constants received in various experiments (Hanks, 1983) confirmed their reasonable consistency from one species to another. However it was noted that the  $m$  constant may vary with varieties within a species. It was pointed out by Fischer and Turner (1978) that  $m$  diverged significantly between  $C_3$  and  $C_4$  species, indicating the evident importance of the photosynthetic pathway in determining  $m$ . Since the basic biochemistry of photosynthesis has not yet been genetically improved within a crop species (Horton, 2000; Parry et al., 2007; Sheehy et al., 2000),  $m$  offers limited scope for significant genetic improvement in biomass at present. The remaining practical and quantitatively effective option for biomass improvement under drought stress is an increase in  $T$  or its seasonal manipulation with respect to plant development and a predictable water supply (see below).

Transpiration efficiency (TE) which is WUE at the leaf level is determined by the delicate interplay between transient photosystem activity, substomatal cavity  $CO_2$  concentration and stomatal activity (Farquhar et al., 1989). Carbon isotope discrimination ( $\delta$ ) measurement in stover or grain has been developed as a method for estimating seasonal plant TE as affected by these transients. Low  $\delta$  is reasonably correlated with high TE (e.g. Hall et al., 1994).  $\delta$  was therefore taken to represent crop WUE (e.g. Condon et al., 2002).

The amount of water transpired per given unit of  $CO_2$  fixation (TE) is an interesting physiological yardstick but our interest here is in maximizing  $CO_2$  fixation under drought stress per plant and per unit land area as an engine of biomass production. Stomatal closure, which is a generally negative response in this respect (not considering survival under severe drought), can be driven by a variety of determinants such as internal leaf  $CO_2$  concentration, cellular solutes, specific ions, pH and ABA produced in the leaf or imported from the root. Higher yielding genotypes of cotton, wheat and rice have greater stomatal conductance and transpiration under drought stress (Araus et al., 2002; Blum et al., 1982; Izanloo et al., 2008; Sanguineti et al., 1999) or well-watered conditions (Fischer et al., 1998; Horie et al., 2006; Lu et al., 1994; Lu and Zeiger, 1994; Reynolds et al., 1994; Shimshi and Ephrat, 1975) as indicated by leaf gas exchange measurements or canopy temperature measurements. Therefore high plant production requires high stomatal conductance over time, to allow greater  $CO_2$  fixation per unit land area, under different conditions. This translates into maximized soil water-use for transpiration. In most cases as will be seen further on, higher  $T$  and high stomatal conductance under drought stress will express lower TE or WUE.

The delicate interplay between transient photosystem activity, substomatal cavity  $CO_2$  concentration and stomatal activity is not the only determinant of plant transpiration and WU at the whole plant and crop level. Since drought stress is the issue here, interest is in plant water-use for stomatal transpiration at given soil

moisture content. Various plant constitutive and stress-adaptive traits have a role in this respect, namely in enhancing an effective use of water (EUW) for transpiration.

Synchronizing growth duration with the expected or the predicted seasonal soil moisture supply is often the first and foremost step in breeding for water-limited environments. Two major considerations are important in this respect: (a) short growth duration dictates moderate water-use and the escape of terminal (reproductive stage) drought stress; and (b) long duration genotypes generally have a greater water-use and larger and deeper root systems that allow deep soil moisture extraction—if indeed deep soil moisture is available (Mitchell et al., 1996).

Leaf permeability is crucial. Leaves can lose water through the cuticle. This non-stomatal “leak” increases crop transpiration without an associated benefit in  $CO_2$  fixation. For example, rice has a highly conductive cuticle while the drought resistant sorghum has a relatively impermeable cuticle. Eliminating this leak by higher epicuticular wax deposition will increase stomatal transpiration (Kerstiens, 1997, 2006). Stomata leakiness at night can also add to water loss without any advantage in  $CO_2$  assimilation. Nocturnal transpiration can be significant (Caird et al., 2007).

Accelerated leaf desiccation and death is a means by which plants reduce water requirement under drought stress. Plant foliage desiccates in a progressive manner from lower (older) leaves to upper (younger) leaves, despite the fact that lower leaves are generally at a better water status than upper leaves. It has been found in sorghum that this strategy reduces transpiration in older and less productive leaves while deviating water-use to younger, fully productive ones (Blum and Arkin, 1984).

An important part of soil moisture available for transpiration is evaporated directly from soil to the atmosphere without taking part in transpiration. For example, up to 40% of the total available soil water was found to be lost by soil evaporation in wheat in Australia (French and Schultz, 1984; Siddique et al., 1990). Soil surface shading by the crop canopy is crucial for reducing this water loss. Reduced soil evaporation by fast vigorous seedling growth is therefore a target in an Australian wheat breeding program (Rebetzke and Richards, 1999) which is actually directed at increasing  $T$ .

The major plant adaptive response to drought at the cellular level which has a proven effect on yield under drought stress is osmotic adjustment (OA) (Blum, 2005). It has been repeatedly demonstrated that OA has two major functions in plant production under drought stress: (a) it enables leaf turgor maintenance for the same leaf water potential thus supporting stomatal conductance under lower leaf water status (e.g. Ali et al., 1999; Sellin, 2001), and (b) it improves root capacity for water uptake (e.g. Chimenti et al., 2006; Tangpremsri et al., 1991).

Therefore the enhancement of biomass production under drought stress can be achieved primarily by maximizing soil water capture while diverting the largest part of the available soil moisture towards stomatal transpiration. This is defined as effective use of water (EUW) and it is the major engine for agronomic or genetic enhancement of crop production under a limited water regime.

There is a wide consensus that the reproductive growth stage is the most sensitive to water deficit. This is well depicted by the classical and widely used example in rice (O’Toole, 1982; Fig. 6). It is also recognized that drought stress at the reproductive stage is the most prevalent problem in rainfed drought prone agriculture, at least simply because in most rainfed ecosystem the crop season’s rains diminish towards flowering and harvest time. Therefore, irrespective of biomass production up to flowering, sustained WU and  $T$  into the reproductive growth stage is crucial for reproductive success (e.g. Merah, 2001; Kato et al., 2008). An effective means of achieving reproductive success under drought

stress is soil moisture capture by deep root system where deep soil moisture is available (e.g. Kirkegaard et al., 2007). ABA production in the shoot or the root under stress may also impede reproductive processes (e.g. Davies and Jones, 1991), but ABA accumulation might be at least partly repressed by higher WU and the resultant improvement of plant water status (Westgate et al., 1996).

Ample information has been developed by carbon isotope discrimination analysis of plants done in order to understand the relationship between TE, WUE and yield under different water-regimes (e.g. Hall et al., 1994). It should be noted here that delta estimates TE at the leaf level which is not crop WUE. However in most studies where delta (TE) was found to be related to yield then the term WUE is used in the report. Following are some major conclusions which serve to reject the notion that high WUE can be equated with drought resistance or the improvement of yield under water-limited conditions (with the exception of crops grown on stored soil moisture (see below).

1. Conflicting results were obtained in various crops under different growing conditions on the association between delta and yield (Hall et al., 1994; Matus et al., 1996; Monneveux et al., 2007; Morgan et al., 1993; Munoz et al., 1998; Ngugi et al., 1994, 1996; Read et al., 1991; Saranga et al., 2004; Sayre et al., 1995; Specht et al., 2001). These range from no relationship between delta and yield to negative or positive relationships, depending on the crop and the environment. Sometimes the relationship was biased by phenology or plant height (or perhaps HI) (e.g. Rebetzke et al., 2008; Sayre et al., 1995). It appears that the association between WUE and plant production as a biologically legitimate variable across species and environments is very elusive. If solid relationships were found they were limited to narrow environmental conditions within a given crop phenology. In this sense WUE in rainfed agriculture is an intangible index. On the other hand its value in economizing the use of expensive irrigation water is well established.
2. Deep or dense root system which would promote soil moisture capture and WU is correlated across genotypes with low WUE (Pinheiro et al., 2005; Kobata et al., 1996) concluded that “the high dry matter production of those rice cultivars known to be drought resistant under field conditions is caused not by high WUE, but by high ability to maintain transpiration, which is supported by deep root systems.”
3. Thus, it is not surprising that favorable genotypic plant water status under drought stress as reflected in measurements of relative water content or canopy temperature is correlated with low WUE across genotypes (Araus et al., 1993; Frank et al., 1997; Read et al., 1991; Zong et al., 2008).
4. Genotypic variation in WUE under limited water regimes is affected more by variation in the denominator (WU) rather than by variation in the nominator (biomass) (Blum, 2005). This has also been determined for TE and stomatal conductance at the single leaf level (e.g. Juenger et al., 2005; Monclus et al., 2006; Monneveux et al., 2006). Hence, selection for high WUE under limited water supply tends to result in a genetic shift towards plant traits that limit crop WU, such as early flowering and smaller leaf area (Martin et al., 1999; Menendez and Hall, 1995; Ngugi et al., 1994; Sayre et al., 1995; White et al., 1990). The successful and widely cited case for dryland wheat grain yield improvement with selection for high WUE (low carbon isotope discrimination) in NSW Australia (Condon et al., 2002) can be explained by the fact that wheat is grown there mainly on stored soil moisture. A major avenue for yield improvement is the control of WU during the earlier part of the growing season in order to avoid lack of soil moisture during reproduction. This was earlier attempted by selection for reduced root xylem diameter (Richards and Passioura, 1989) and it can also be

achieved by reduced leaf area and growth duration as done in the past in sorghum grown under stored soil moisture conditions (Blum, 1970, 1972; Blum and Naveh, 1976). Such plants that allow optimized seasonal distribution of soil moisture use express high WUE for grain yield due to their relative moderate WU and high HI. The same genetic materials, selected for high WUE were not successful in Western Australia where rainfed wheat does not grow on stored soil moisture (Fig. 6 in Condon et al., 2002).

5. Considering all of the above (1 through 4), it is not surprising that drought resistance was found to be associated with low WUE when analyzed by delta under limited water supply (e.g. Araus et al., 2003; Morgan et al., 1993; Ngugi et al., 1994; Solomon and Labuschagne, 2004). A drought resistant *Coffea canephora* clone had relatively lower WUE than a drought susceptible one, where resistance was associated with deeper roots and presumably greater WU (Pinheiro et al., 2005).
6. Finally, crop WUE has long been known to increase with increasing drought stress and reduced water supply (e.g. Meyers et al., 1984). This has been more recently confirmed with delta analysis (Craufurd et al., 1999; Ismail et al., 1994; Li et al., 2000; Peuke et al., 2006). It corresponds well with the fact that plant water deficit result in high WUE (#3 above). Assume therefore that two different cultivars are planted side by side and exposed to drought stress. If the one with higher WUE is selected it will most likely to be the one relatively more stressed and at a lower plant water status, namely the drought susceptible one.

Therefore, for all practical purposes plant breeders targeting water-limited environments should consider skipping the use and reference to WUE and consider plant constitutive and adaptive traits which drive the effective use of water (EUW) and the resultant dehydration avoidance as major traits for yield improvement in drought prone environments (with one exception per #4 above). This discussion does not refer to very shallow soils with very limited soil water holding capacity. These extremely difficult conditions require another discussion on plant survival and recovery and not plant production.

In conclusion, crop WUE as estimated under rainfed conditions by carbon isotope discrimination analysis or any other method is an elusive ratio. Reynolds and Tuberosa (2008) concluded in tune with the original expression of Passioura (1996) that “water uptake (WU), water-use efficiency (WUE), and harvest index (HI) are drivers of yield.” Indeed WU and HI are drivers of yield but I suggest that WUE is just a passenger. Whereas HI (in terms of assimilate partitioning and reproductive success under drought stress) is also largely influenced by WU and plant water status, it can be concluded that WU alone is the main (not the exclusive) driver of yield under drought stress.

Therefore, it is not a coincidence that EUW is an inverse acronym of WUE whereas very often high WUE is achieved at the expense of reduced EUW, and *vice versa*.

## References

- Ali, M., Jensen, C.R., Mogensen, V.O., Andersen, M.N., Henson, I.E., 1999. Root signaling and osmotic adjustment during intermittent soil drying sustain grain yield of field grown wheat. *Field Crops Res.* 62, 35–52.
- Araus, J.L., Reynolds, M.P., Acevedo, E., 1993. Leaf posture, grain yield, growth, leaf structure, and carbon isotope discrimination in wheat. *Crop Sci.* 33, 1273–1279.
- Araus, J.L., Slafer, G.A., Reynolds, M.P., Royo, C., 2002. Plant breeding and water stress in C3 cereals: what to breed for? *Ann. Bot.* 89, 925–940.
- Araus, J.L., Villegas, D., Aparicio, N., García Del Moral, L.F., El Hani, S., Rharabti, Y., Ferrio, J.P., Royo, C., 2003. Environmental factors determining carbon isotope discrimination and yield in durum wheat under Mediterranean conditions. *Crop Sci.* 43, 170–180.
- Blum, A., 1970. Effects of plant density and growth duration on sorghum yield under limited water supply. *Agron. J.* 62, 333–336.

- Blum, A., 1972. Effect of planting date on water-use and its efficiency in dryland grain sorghum. *Agron. J.* 64, 775–778.
- Blum, A., 2005. Drought resistance, water-use efficiency, and yield potential—are they compatible, dissonant, or mutually exclusive? *Aust. J. Agric. Res.* 56, 1159–1168.
- Blum, A., Arkin, G.F., 1984. Sorghum root growth and water-use as affected by water supply and watering duration. *Field Crops Res.* 9, 131–142.
- Blum, A., Mayer, J., Gozlan, G., 1982. Infrared thermal sensing of plant canopies as a screening technique for dehydration avoidance in wheat. *Field Crops Res.* 5, 137–146.
- Blum, A., Naveh, M., 1976. Improved water-use efficiency by promoted plant competition in dryland sorghum. *Agron. J.* 68, 111–116.
- Caird, M.A., Richards, J.H., Hsiao, T.C., 2007. Significant transpirational water loss occurs throughout the night in field-grown tomato. *Funct. Plant Biol.* 34, 172–177.
- Chimentì, C.A., Marcantonio, M., Hall, A.J., 2006. Divergent selection for osmotic adjustment results in improved drought tolerance in maize (*Zea mays* L.) in both early growth and flowering phases. *Field Crops Res.* 95, 305–315.
- Condon, A.G., Richards, R.A., Rebetzke, G.J., Farquhar, G.D., 2002. Improving intrinsic water-use efficiency and crop yield. *Crop Sci.* 42, 122–131.
- Craufurd, P.Q., Wheeler, T.R., Ellis, R.H., Summerfield, R.J., Williams, J.H., 1999. Effect of temperature and water deficit on water-use efficiency, carbon isotope discrimination and specific leaf area in peanut. *Crop Sci.* 39, 136–142.
- Davies, W.J., Jones, H.G., 1991. *Abscisic Acid: Physiology and Biochemistry*. Bios Scientific Publishers, London, p. 266.
- de Wit, C.T., 1958. *Transpiration and Crop Yields*, vol. 64.6. Versl. Landbouwk. Onderz., Institute of Biological and Chemical Research on Field Crops and Herbage, Wageningen, The Netherlands, pp. 24.
- Donald, C.M., Hamblin, J., 1976. The biological yield and harvest index of cereals as agronomic and plant breeding criteria. *Adv. Agron.* 28, 361–405.
- Farquhar, G.D., Ehleringer, J.R., Hubick, K., 1989. Carbon isotope discrimination and photosynthesis. *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 40, 503–537.
- Fischer, R.A., Turner, N.C., 1978. Plant productivity in the arid and semiarid zones. *Ann. Rev. Plant Physiol.* 29, 277–317.
- Fischer, R.A., Rees, D., Sayre, K.D., Lu, Z.M., Condon, A.G., Saavedra, A.L., 1998. Wheat yield progress associated with higher stomatal conductance and photosynthetic rate, and cooler canopies. *Crop Sci.* 38, 1467–1475.
- Frank, A.B., Ray, I.M., Berdahl, J.D., Karn, J.F., 1997. Carbon isotope discrimination, ash, and canopy temperature in three wheatgrass species. *Crop Sci.* 7, 1573–1576.
- French, R.J., Schultz, J.E., 1984. Water use efficiency of wheat in a Mediterranean-type environment. I. The relation between yield water use and climate. *Aust. J. Agric. Res.* 35, 743–764.
- Hall, A.E., Richards, R.A., Condon, A.G., Wright, G.C., Farquhar, G.D., 1994. Carbon isotope discrimination and plant breeding. *Plant Breed. Rev.* 12, 91–113.
- Hanks, K.J., 1983. Yield and water-use relationships, an overview. In: Taylor, H.M., Jordan, W.R., Sinclair, T.R. (Eds.), *Limitations to Efficient Water Use in Crop Production*. American Society of Agronomy, Madison, WI (USA), pp. 393–410.
- Horie, T., Matsuura, S., Takai, T., Kuwasaki, K., Ohsumi, A., Shiraiwa, T., 2006. Genotypic difference in canopy diffusive conductance measured by a new remote-sensing method and its association with the difference in rice yield potential. *Plant Cell Environ.* 29, 653–660.
- Horton, P., 2000. Prospects for crop improvement through the genetic manipulation of photosynthesis, morphological and biochemical aspects of light capture. *J. Exp. Bot.* 51, 475–485.
- Ismail, A.M., Hall, A.E., Bray, E.A., 1994. Drought and pot size effects on transpiration efficiency and carbon isotope discrimination of cowpea accessions and hybrids. *Aust. J. Plant Physiol.* 21, 23–35.
- Izanloo, A., Condon, A.G., Langridge, P., Tester, M., Schnurbusch, T., 2008. Different mechanisms of adaptation to cyclic water stress in two South Australian bread wheat cultivars. *J. Exp. Bot.* 59, 3327–3346.
- Juenger, T.E., McKay, J.K., Hausmann, N., Keurentjes, J.J.B., Sen, S., Stowe, K.A., Dawson, T.E., Simms, E.L., Richards, J.H., 2005. Identification and characterization of QTL underlying whole-plant physiology in *Arabidopsis thaliana*, <sup>13</sup>C, stomatal conductance and transpiration efficiency. *Plant Cell Environ.* 28, 697–708.
- Kato, Y., Kamoshita, A., Yamagishi, J., 2008. Preflowering abortion reduces spikelet number in upland rice (*Oryza sativa* L.) under water stress. *Crop Sci.* 48, 2389–2395.
- Kerstiens, G., 1997. In vivo manipulation of cuticular water permeance and its effect on stomatal response to air humidity. *New Phytol.* 137, 473–480.
- Kerstiens, G., 2006. Water transport in plant cuticles, an update. *J. Exp. Bot.* 57, 2493–2499.
- Kijne, J.W., Barker, R., Molden, D.J. (Eds.), 2003. *Water Productivity in Agriculture: Limits and Opportunities for Improvement*. CABI, UK, p. 332.
- Kirkegaard, J.A., Lilley, J.M., Howe, G.N., Graham, J.M., 2007. Impact of subsoil water use on wheat yield. *Aust. J. Agric. Res.* 58, 303–315.
- Kobata, T., Okuno, T., Yamamoto, T., 1996. Contributions of capacity for soil water extraction and water use efficiency to maintenance of dry matter production in rice subjected to drought. *Jpn. J. Crop Sci.* 65, 652–662.
- Li, C.Y., Berninger, F., Koskela, J., Sonninen, E., 2000. Drought responses of *Eucalyptus microtheca* provenances depend on seasonality of rainfall in their place of origin. *Aust. J. Plant Physiol.* 27, 231–238.
- Lu, Z.M., Zeiger, E., 1994. Selection for higher yields and heat resistance in pima cotton has caused genetically determined changes in stomatal conductances. *Physiol. Plant.* 92, 273–278.
- Lu, Z.M., Radin, J.W., Turcotte, E.L., Percy, R., Zeiger, E., 1994. High yields in advanced lines of pima cotton are associated with higher stomatal conductance, reduced leaf area and lower leaf temperature. *Physiol. Plant.* 92, 266–272.
- Martin, B., Tauer, C.G., Lin, R.K., 1999. Carbon isotope discrimination as a tool to improve water-use efficiency in tomato. *Crop Sci.* 39, 1775–1783.
- Matus, A., Slinkard, A.E., Vankessel, C., 1996. Carbon isotope discrimination and indirect selection for transpiration efficiency at flowering in lentil (*Lens culinaris medikus*), spring bread wheat (*Triticum aestivum* L.) durum wheat (*T. turgidum* L.) and canola (*Brassica napus* L.). *Euphytica* 87, 141–151.
- Menendez, C.M., Hall, W.E., 1995. Heritability of carbon isotope discrimination and correlations with earliness in cowpea. *Crop Sci.* 35, 673–678 3032.
- Merah, O., 2001. Potential importance of water status traits for durum wheat improvement under Mediterranean conditions. *J. Agric. Sci.* 137, 139–145.
- Meyers, R.J.K., Foale, M.A., Done, A.A., 1984. Response of grain sorghum to varying irrigation frequency in the Ord irrigation area. II. Evapotranspiration water-use efficiency. *Aust. J. Agric. Res.* 35, 31–42.
- Mitchell, J.H., Fukai, S., Cooper, M., 1996. Influence of phenology on grain yield variation among barley cultivars grown under terminal drought. *Aust. J. Agric. Res.* 47, 757–774.
- Monclus, R., Dreyer, E., Villar, M., Delmotte, F.M., Delay, D., Petit, J.M., Barbaroux, C., Thiec, D., Bréchet, C., Brignolas, F., 2006. Impact of drought on productivity and water use efficiency in 29 genotypes of *Populus deltoides* × *Populus nigra*. *New Phytol.* 169, 765–777.
- Monneveux, P., Rekika, D., Acevedo, E., Merah, O., 2006. Effect of drought on leaf gas exchange, carbon isotope discrimination, transpiration efficiency and productivity in field grown durum wheat genotypes. *Plant Sci.* 170, 867–872.
- Monneveux, P., Sheshshayee, M.S., Akhter, J., Ribaut, J.M., 2007. Using carbon isotope discrimination to select maize (*Zea mays* L.) inbred lines and hybrids for drought tolerance. *Plant Sci.* 173, 390–396.
- Morgan, J.A., Lecain, D.R., Mccaig, T.N., Quick, J.S., 1993. Gas exchange, carbon isotope discrimination, and productivity in winter wheat. *Crop Sci.* 33, 178–186.
- Munoz, P., Voltas, J., Araus, J.L., Igarua, E., Romagosa, I., 1998. Changes over time in the adaptation of barley releases in north-eastern Spain. *Plant Breed.* 117, 531–535.
- Ngugi, E.C.K., Austin, R.B., Galwey, N.W., Hall, M.A., 1996. Associations between grain yield and carbon isotope discrimination in cowpea. *Eur. J. Agron.* 5, 9–17.
- Ngugi, E.C.K., Galwey, N.W., Austin, R.B., 1994. Genotype × environment interaction in carbon isotope discrimination and seed yield in cowpea (*Vigna unguiculata* L. Walp.). *Euphytica* 73, 213–224.
- O'Toole, J.C., 1982. Adaptation of rice to drought-prone environments. In: *Drought Resistance in Crops With Emphasis on Rice*, IRR, International Rice Research Institute, Los Banos, Philippines, pp. 195–213.
- Parry, M.A.J., Madgwick, P.J., Carvahlo, J.F.C., Andralojo, P.J., 2007. Prospects for increasing photosynthesis by overcoming the limitations of Rubisco. *J. Agric. Sci.* 145, 31–43.
- Passioura, J.B., 1996. Drought and drought tolerance. *Plant Growth Regul.* 20, 79–83.
- Passioura, J.B., 2006. Increasing crop productivity when water is scarce—from breeding to field management. *Agric. Water Manage.* 80, 176–196.
- Peuke, A.D., Gessler, A., Rennenberg, H., 2006. The effect of drought on C and N stable isotopes in different fractions of leaves, stems and roots of sensitive and tolerant beech ecotypes. *Plant Cell Environ.* 29, 823–835.
- Pinheiro, H.A., Damatta, F.M., Chaves, A.R.M., Loureiro, M.E., 2005. Drought tolerance is associated with rooting depth and stomatal control of water use in clones of *Coffea canephora*. *Ann. Bot.* 96, 101–108.
- Read, J.J., Johnson, D.A., Asay, K.H., Tieszen, L.L., 1991. Carbon isotope discrimination, gas exchange, and water-use efficiency in crested wheatgrass clones. *Crop Sci.* 31, 1203–1208.
- Rebetzke, G.J., Condon, A.G., Farquhar, G.D., Appels, R., Richards, R.A., 2008. Quantitative trait loci for carbon isotope discrimination are repeatable across environments and wheat mapping populations. *Theor. Appl. Gen.* 118, 123–137.
- Rebetzke, G.J., Richards, R.A., 1999. Genetic improvement of early vigour in wheat. *Aust. J. Agric. Res.* 50, 291–301.
- Reynolds, M.P., Balota, M., Delgado, M.I.B., Amani, I., Fischer, R.A., 1994. Physiological and morphological traits associated with spring wheat yield under hot, irrigated conditions. *Aust. J. Plant Physiol.* 21, 717–730.
- Reynolds, M., Tuberosa, R., 2008. Translational research impacting on crop productivity in drought-prone environments. *Curr. Opin. Plant Biol.* 11, 171–179.
- Richards, R.A., Passioura, J.B., 1989. A breeding program to reduce the diameter of the major xylem vessel in the seminal roots of wheat and its effect on grain yield in rain-fed environments. *Aust. J. Agric. Res.* 40, 943–950.
- Sanguinetti, M.C., Tuberosa, R., Landi, P., Salvi, S., Maccaferri, M., Casarini, E., Conti, S., 1999. QTL analysis of drought related traits and grain yield in relation to genetic variation for leaf abscisic acid concentration in field-grown maize. *J. Exp. Bot.* 50, 1289–1297.
- Saranga, Y., Jiang, C.X., Wright, R.J., Yakir, D., Paterson, A.H., 2004. Genetic dissection of cotton physiological responses to arid conditions and their inter-relationships with productivity. *Plant Cell Environ.* 27, 263–277.
- Sayre, K.D., Acevedo, E., Austin, R.B., 1995. Carbon isotope discrimination and grain yield for three bread wheat germplasm groups grown at different levels of water stress. *Field Crop Res.* 41, 45–54.
- Sellin, A., 2001. Hydraulic and stomatal adjustment of Norway spruce trees to environmental stress. *Tree Physiol.* 21, 879–888.

- Sheehy, J.E., Mitchell, P.L., Hardy, B. (Eds.), 2000. Redesigning Rice Photosynthesis to Increase Yield. Elsevier Science, Amsterdam (The Netherlands), p. 300.
- Shimshi, D., Ephrat, J., 1975. Stomatal behavior of wheat cultivars in relation to their transpiration, photosynthesis and yield. *Agron. J.* 67, 326–331.
- Siddique, K.H.M., Tennant, D., Perry, M.W., Belford, R.K., 1990. Water use and water use efficiency of old and modern wheat cultivars in a Mediterranean-type environment. *Aust. J. Agric. Res.* 41, 431–447.
- Solomon, K.F., Labuschagne, M.T., 2004. Variation in water use and transpiration efficiency among durum wheat genotypes grown under moisture stress and non-stress conditions. *J. Agric. Sci.* 141, 31–41.
- Specht, J.E., Chase, K., Macrander, M., Graef, G.L., Chung, J., Markwell, J.P., Germann, M., Orf, J.H., Lark, K.G., 2001. Soybean response to water; a QTL analysis of drought tolerance. *Crop Sci.* 41, 493–509.
- Tangpremsri, T., Fukai, S., Fischer, K.S., Henzell, R.G., 1991. Genotypic variation in osmotic adjustment in grain sorghum. 2. Relation with some growth attributes. *Aust. J. Agric. Res.* 42, 759–767.
- Westgate, M.E., Passioura, J.B., Munns, R., 1996. Water status and ABA content of floral organs in drought-stressed wheat. *Aust. J. Plant Physiol.* 23, 763–772.
- White, J.W., Castillo, J.A., Ehleringer, J., 1990. Associations between productivity, root growth and carbon isotope discrimination in *Phaseolus vulgaris* under water deficit. *Aust. J. Plant Physiol.* 17, 189–198.
- Zong, L.Z., Liang, S., Xu, X., Li, S.H., Jing, J.H., Monneveux, P., 2008. Relationships between carbon isotope discrimination and leaf morpho-physiological traits in spring-planted spring wheat under drought and salinity stress in Northern China. *Aust. J. Agric. Res.* 59, 941–949.